



<http://www.biodiversitylibrary.org>

Entomological news.

[Philadelphia]American Entomological Society, 1925-
<http://www.biodiversitylibrary.org/bibliography/2356>

v. 104-105 1993: <http://www.biodiversitylibrary.org/item/20581>
Page(s): Page 277, Page 278, Page 279

Contributed by: Smithsonian Institution Libraries
Sponsored by: Smithsonian

Generated 13 April 2010 11:35 AM
<http://www.biodiversitylibrary.org/pdf2/002825500020581>

This page intentionally left blank.

NOTE ON THE DISTRIBUTION OF THE IMMIGRANT BARK BEETLE, *HYLASTES OPACUS*, IN NORTH AMERICA (COLEOPTERA: SCOLYTIDAE)¹

Robert J. Rabaglia², J. F. Cavey³

ABSTRACT: The Palearctic bark beetle, *Hylastes opacus*, was first recorded for North America in 1989 from a single location near Long Island, New York. This paper reports *H. opacus* from Maine, New Hampshire, Vermont, West Virginia, and central New York. This black pine bark beetle was detected in Maine and Vermont in *Ips* pheromone-baited Lindgren funnel traps in 1993, in alpha-pinene baited traps in New Hampshire in 1994, and an ethanol baited trap in West Virginia in 1994. Additional specimens were identified from an earlier collection in New York. These records extend the known range of *H. opacus* in North America from New York to northern New England and south to West Virginia..

The palearctic bark beetle, *Hylstes opacus* Erichson occurs in the "pine belts" of Asia and Europe (China, Japan, Korea, Austria, Belgium, Bulgaria, Denmark, England, Finland, France, Germany, Greece, Norway, Poland, Sweden, Switzerland and USSR) (Wood and Bright 1992). The species has also become established in pine plantations of South Africa (Wood 1992).

Wood (1992) reported the first North American record of *H. opacus*, from a series collected on Fisher's Island, Suffolk County, New York, in May, 1989 by T. W. Phillips. Based on the circumstances of the collection, Wood (1992) concluded that the *H. opacus* was established at this site.

The Fisher's island collection consisted of 12 specimens in a pheromone trap. The island is devoid of native pines, with only one potential host for *H. opacus* – ornamental plantings of Japanese black pine, *Pinus thunbergiana* Franco. (T.W. Phillips, pers. commun., Feb. 1994).

In 1993, we detected specimens of *H. opacus* from Vermont and Maine while screening trap samples from a Cooperative Agricultural Pest Survey (CAPS) for exotic bark beetles in the Northeastern United States. These records extend the known geographic range for *H. opacus* in North America to the northeast from New York to New Engand (Figure 1). Specific collection data area as follows:

Maine: Waldo Co., Lincolnville, 6/V/1993, R. Mack Collector, four specimens and **Vermont:** Washington Co., Plainfield, 5/VI/1993, M. Michaelis Collector and Washington Co., Graniteville, 3/VI/1993, M. Michaelis Collector, single specimens.

¹ Received June 9, 1994. Accepted July 12, 1994.

² Maryland Department of Agriculture, 50 Harry S. Truman Pkwy, Annapolis, MD 21401.

³ U.S. Department of Agriculture, Animal Health and Plant Inspection Service, Plant Protection and Quarantine (USDA-APHIS-PPQ), Federal Bldg, Room 625, 6506 Belcrest Rd, Hyattsville, MD 20782.

All collections were from Lindgren funnel traps baited with Ipslure (ipsdienol, methylbutanol and cis-verbenol). Specimens were identified using Balachowsky (1949) and by comparison to European material in the Eggers Collection at the U.S. National Museum (USNM).

After reviewing the new England specimens, the senior author identified a series of *H. opacus* he collected from a cut stump of *Pinus sylvestris* L. in **New York**: Oneida Co., Woodgate, 27/IV/1987. This Oneida County record is additional to 1993 collections of *H. opacus* reported from 22 New York counties by Hoebeke 1994.




Figure 1. Collection records of *Hylastes opacus*: Star – first collection in North America (Wood 92); Closed circles – new state and county records reported in this paper.

During the review period for the manuscript we learned of the following records of *H. opacus*: from **New Hampshire**: Cheshire Co., Keene, 14-22/IV/1994, J.S. Weaver and S. Longsjoe collectors, in pine shoot beetle trap baited with alpha-pinene (identified by Dr. Donald E. Bright, Agriculture Canada, Ottawa, Canada). **West Virginia**: Wood Co., Parkersburg, 1-8/VI/1994, R. A. Meyers collector, in ethanol-baited Lindgren funnel trap (identified by Dr. Steven L. Wood, Brigham Young University, Provo, UT).

One specimen from Maine and one from New York are deposited in the USNM Insect Collection.

ACKNOWLEDGMENTS

We thank Natalia J. Vandenberg, USDA, Agricultural Research Service, Systematic Entomology Laboratory (ARS-SEL), Washington D.C. and Donald M. Anderson, USDA-ARS-SEL (retired) for assistance in using the USNM collections and library; Richard Mack, Maine Pest Survey Coordinator, Maine Cooperative Extension Service, Orono, and Mark Michaelis, USDA-APHIS-PPQ, Berlin, Vermont, for collecting the specimens; John S. Weaver, Department of Entomology, University of New Hampshire, Durham, for contributing the New Hampshire state record; R. A. Meyers, West Virginia Department of Agriculture, for contributing the West Virginia record; and E. Richard Hoebeke, Cornell University, Ithaca, New York and Charles L. Staines, Maryland Department of Agriculture, Annapolis for reviewing an earlier draft of this note.

LITERATURE CITED

- Balachowsky, A.** 1949. Coléoptères Scolytides. Faune de France. 50:1-320.
- Hoebeke, E.R.** 1994. New records of immigrant bark beetles (Coleoptera:Scolytidae) in New York: Attraction of conifer infesting species to ethanol-baited trap logs. Entomol. News 105(5): 267-276.
- Wood, S.L.** 1992. Nomenclatural changes and new species in Platypodidae and Scolytidae (Coleoptera), Part II, Great Basin Nat. 52(1):78-88.
- Wood, S.L. and D.E. Bright, Jr.** 1992. A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index. Volume A (pp 1-833), volume B (pp 835-1553). Great Basin Nat. Mem. 13.

The following text is generated from uncorrected OCR.

[Begin Page: Page 277]

Vol. 105, No. 5, November & December, 1994 277

NOTE ON THE DISTRIBUTION OF THE IMMIGRANT

BARK BEETLE, *HYLASTES OPACUS*, IN NORTH

AMERICA (COLEOPTERA: SCOLYTIDAE) 1

Robert J. Rabaglia 2 , J. F. Cavey 3

ABSTRACT: The Palearctic bark beetle, *Hylastes opacus*, was first recorded for North America in 1989 from a single location near Long Island, New York. This paper reports *H. opacus* from Maine, New Hampshire, Vermont, West Virginia, and central New York. This black pine bark beetle was detected in Maine and Vermont in Ips pheromone-baited Lindgren funnel traps in 1993, in alpha-pinene baited traps in New Hampshire in 1994, and an ethanol baited trap in West Virginia in 1994. Additional specimens were identified from an earlier collection in New York. These records extend the known range of *H. opacus* in North America from New York to northern New England and south to West Virginia..

The palearctic bark beetle, *Hylstes opacus* Erichson occurs in the "pine belts" of Asia and Europe (China, Japan, Korea, Austria, Belgium, Bulgaria, Denmark, England, Finland, France, Germany, Greece, Norway, Poland, Sweden, Switzerland and USSR) (Wood and Bright 1992). The species has also become established in pine plantations of South Africa (Wood 1992).

Wood (1992) reported the first North American record of *H. opacus*, from a series collected on Fisher's Island, Suffolk County, New York, in May, 1989 by T. W. Phillips. Based on the circumstances of the collection, Wood (1992) concluded that the *H. opacus* was established at this site.

The Fisher's island collection consisted of 12 specimens in a pheromone trap. The island is devoid of native pines, with only one potential host for *H. opacus* - ornamental plantings of Japanese black pine, *Pinus thunbergiana* Franco. (T.W. Phillips, pers. commun., Feb. 1994).

In 1993, we detected specimens of *H. opacus* from Vermont and Maine while screening trap samples from a Cooperative Agricultural Pest Survey (CAPS) for exotic bark beetles in the Northeastern United States. These records extend the known geographic range for *H. opacus* in North America to the northeast from New York to New England (Figure 1). Specific collection data area as follows:

Maine: Waldo Co., Lincolnville, 6/V/1993, R. Mack Collector, four specimens and Vermont: Washington Co., Plainfield, 5/VI/1993, M. Michaelis Collector and Washington Co., Graniteville, 3/VI/1993, M. Michaelis Collector, single specimens.

1 Received June 9, 1994. Accepted July 12, 1994

2 Maryland Department of Agriculture, 50 Harry S. Truman Pkwy, Annapolis, MD 21401.

3 U.S. Department of Agriculture, Animal Health and Plant Inspection Service, Plant Protection and Quarantine (USDA-APHIS-PPQ), Federal Bldg, Room 625, 6506 Belcrest Rd, Hyattsville,

MD 20782.

ENT. NEWS 105(5): 277-279, November & December 1994

[Begin Page: Page 278]

278

ENTOMOLOGICAL NEWS

All collections were from Lindgren funnel traps baited with Ipslure (ipsdienol, methylbutanol and cis-verbenol). Specimens were identified using Balachowsky (1949) and by comparison to European material in the Eggers Collection at the U.S. National Museum (USNM).

After reviewing the new England specimens, the senior author identified a series of *H. opacus* he collected from a cut stump of *Pinus sylvestris* L. in New York: Oneida Co., Woodgate, 27/IV/1987. This Oneida County record is additional to 1993 collections of *H. opacus* reported from 22 New York counties by Hoebeke 1994.

Figure 1. Collection records of *Hylastes opacus*: Star - first collection in North America (Wood 92); Closed circles - new state and county records reported in this paper.

[Begin Page: Page 279]

During the review period for the manuscript we learned of the following records of *H. opacus*: from New Hampshire: Cheshire Co., Keene, 14-22/IV/1994, J.S. Weaver and S. Longsjoe collectors, in pine shoot beetle trap baited with alpha-pinene (identified by Dr. Donald E. Bright, Agriculture Canada, Ottawa, Canada). West Virginia: Wood Co., Parkersburg, 1-8/VI/1994, R. A. Meyers collector, in ethanol-baited Lindgren funnel trap (identified by Dr. Steven L. Wood, Brigham Young University, Provo, UT).

One specimen from Maine and one from New York are deposited in the USNM Insect Collection.

ACKNOWLEDGMENTS

We thank Natalia J. Vandenberg, USDA, Agricultural Research Service, Systematic Entomology Laboratory (ARS-SEL), Washington D.C. and Donald M. Anderson, USDA-ARS-SEL (retired) for assistance in using the USNM collections and library; Richard Mack, Maine Pest Survey Coordinator, Maine Cooperative Extension Service, Orono, and Mark Michaelis, USDA-APHIS-PPQ, Berlin, Vermont, for collecting the specimens; John S. Weaver, Department of Entomology, University of New Hampshire, Durham, for contributing the New Hampshire state record; R. A. Meyers, West Virginia Department of Agriculture, for contributing the West Virginia record; and E. Richard Hoebeke, Cornell University, Ithaca, New York and Charles L. Staines, Maryland Department of Agriculture, Annapolis for reviewing an earlier draft of this note.

LITERATURE CITED

Balachowsky, A. 1949. Coleopteres Scolytides. Faune de France. 50:1-320.

Hoebeke, E.R. 1994. New records of immigrant bark beetles (Coleoptera:Scolytidae) in New York: Attraction of conifer infesting species to ethanol-baited trap logs. Entomol. News 105(5): 267-276.

Wood, S.L. 1992. Nomenclatural changes and new species in Platypodidae and Scolytidae (Coleoptera), Part II, Great Basin Nat. 52(1):78-88.

Wood, S.L. and D.E. Bright, Jr. 1992. A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index. Volume A (pp 1-833), volume B (pp 835-1553). Great Basin Nat. Mem. 13.